


Geophysical surveys for the characterization of glacial and periglacial environment

(Proposer: Jacopo Boaga)

The characterization of glacial and periglacial environments is crucial for several reasons, from the consequences on potential slope instability to the monitoring of the climate change effects in the high-altitude alpine environment. Permafrost is the well-known ground layer with a temperature remaining at or below 0°C for at least two consecutive years. Permafrost interests one quarter of the Northern Hemisphere and 17% of the entire Earth (Biskaborn et al 2019), and is intensively studied from decades in the polar regions and in the high mountain environments (Phillips et al 2009). Geophysical prospecting can usefully help to extend the characterization of high mountains areas, increasing the characterization of the permafrost zones (Hauck 2001; Scott et al. 1990, Boaga et al. 2020, Pavoni et al. 2021) and the study of rock glacier environment. The PhD project regards in particular the use of electrical resistivity tomography (ERT), Frequency Domain Electro-magnetic techniques (FDEM) radar (GPR) and refraction seismic (RST) in several sites of the Alps already monitored with different techniques. The aim is to test limit and applicability of the methods and in particular test the use of contactless geophysics to characterize the permafrost active layer (in terms of layer thickness and lateral continuity) and the rock glacier structures. The work will involve national and international collaboration with the WSL - SLF institute of Davos (Switzerland), the Province of Trento (Italy) and CNR -ISP (Italy).



Available funds: FINA Project ‘Geofisica applicata all’ambiente e al territorio

References

- M Pavoni, F Sirch, J Boaga, Electrical and Electromagnetic Geophysical Prospecting for the Monitoring of Rock Glaciers in the Dolomites, Northeast Italy. Sensors 21 (4), 1294
- Boaga, J.; Phillips, M.; Noetzli, J.; Haberkorn, A.; Kenner, R.; Bast, A. 2020. A Comparison of Frequency Domain Electro-Magnetometry, Electrical Resistivity Tomography and Borehole Temperatures to Assess the Presence of Ice in a Rock Glacier. DOI:10.3389/feart.2020.586430. In Frontiers in Earth Science - ISSN:2296-6463 vol. 8
- Biskaborn, B.K., Smith, S. L., Noetzli J.,Matthes H.,Vieira G., Streletsiky D. A.,Schoeneich P.,Romanovsky V. E., Lewkowicz A. G., Abramov A., Allard M., Boike J.,Cable W. L., Christiansen H. H., Delaloye R.,Diekmann B., Drozdov D., Etzelmüller B., Grosse G.,Guglielmin M., Ingeman-Nielsen T., Isaksen K., Ishikawa M., Johansson M., Johannsson H., Joo A., Kaverin D., Kholodov A., Konstantinov P., Kröger T., Lambiel C.,Lanckman JP., Luo D.,Malkova G.,Meiklejohn I.,Moskalenko N.,Oliva M., Phillips M., Ramos M., Sannel A. B. K., Sergeev D., Seybold C., Skryabin P., Vasiliev A., Wu Q., Yoshikawa K., Zheleznyak M.,Lantuit H. 2019 . Permafrost is warming at a global scale. Nature Communications, volume 10 doi.org/10.1038/s41467-018-08240
- Hauck, C., Guglielmin, M., Isaksen, K. and Yonder l'vliihll, D., 2001. Applicability of frequency-domain and time-domain electromagnetic methods for mountain permafrost studies. Permafrost and Periglacial Processes, 12(1), 39-52.
- Phillips, M., Mutter, E.Z., Kern-Luetschg, M., Lehning M., 2009. Rapid degradation of ground ice in a ventilated talus slope: Flüela Pass, Swiss Alps. Permafrost and Periglacial Processes 20(1), pp. 1-14

- Scott, W., Sellmann, P. and Hunter, J., 1990. Geophysics in the study of permafrost. In: Geotechnical and Environmental Geophysics, (ed. S.Ward), pp. 355- 384. Soc. of Expl. Geoph., Thlsa.